

A close-up photograph of a small, light brown snail crawling on a large, translucent green leaf. The leaf's veins are clearly visible, and the background is a soft-focus green.

L7 - Classical Mechanics 3

Molecular mechanics: dynamics

Born-Oppenheimer Molecular Dynamics (BOMD)

Second Newton's law:


$$M_\alpha \frac{d^2 \mathbf{R}_\alpha}{dt^2} = -\nabla_\alpha E_K(\mathbf{R})$$

Solve it numerically, for instance, with Velocity Verlet:

$$\mathbf{R}_\alpha(t + \Delta t) = \mathbf{R}_\alpha(t) + \mathbf{v}_\alpha(t)\Delta t + \frac{1}{2}\mathbf{a}_\alpha(t)\Delta t^2$$

$$\mathbf{v}_\alpha(t + \Delta t) = \mathbf{v}_\alpha(t) + \frac{1}{2}(\mathbf{a}_\alpha(t) + \mathbf{a}_\alpha(t + \Delta t))\Delta t$$

Classical EOM



The nuclei move on the potential energy surface $E_K(\mathbf{R})$ of electronic state K

$$(T_{elec}(\mathbf{r}) + V(\mathbf{r}, \mathbf{R}))\varphi_K(\mathbf{r}; \mathbf{R}) = E_K(\mathbf{R})\varphi_K(\mathbf{r}; \mathbf{R})$$

How do we get this surface?

- Precompute
 - ✓ Fitting
 - ✓ Modeling
- Compute as needed

PES type		Pros	Cons
Precomputed	Fitted	<ul style="list-style-type: none"> • May consider nonlocal effects (tunneling, ...) 	<ul style="list-style-type: none"> • Curse of dimensionality • Difficult procedure • Expensive preparation
	Modeled	<ul style="list-style-type: none"> • May consider nonlocal effects (tunneling, ...) 	<ul style="list-style-type: none"> • Lack of functional flexibility • Expensive preparation
Computed as needed		<ul style="list-style-type: none"> • Offer QM flexibility (easy to change methods) 	<ul style="list-style-type: none"> • Miss nonlocal effects • Expensive dynamics

Fitted precomputed PES


A collection of fitted potential energy surfaces is available at
comp.chem.umn.edu/potlib/form.cgi

Non-fitted datasets:

- MD17 & MD22: www.sgdml.org/#datasets
- QM-22: github.com/jmbowma/QM-22
- WS22: zenodo.org/record/6985377

Machine learning is reviving the field

1. Build a dataset of energies and energy gradients (100 k points)
2. Train a machine (kernel, NN) to predict energies and gradients for unseen geometries



- MD17 Database
- Energy + Force
- $N_{\text{train}} = 1\text{k}; N_{\text{model}} = 20; N_{\text{test}} = 20\text{k}$

Model PES

Model Hamiltonians assume that the PES has some specific shape.

Thus, the amount of computed data is smaller than in fitted surfaces.

They are needed to parameterize the model.

Examples of model Hamiltonians:


- Spin-Boson Hamiltonian
- Linear Vibronic Coupling
- Molecular mechanics force fields

Example: Spin-Boson model

$$V_a(\mathbf{R}) = \varepsilon_0 + \frac{1}{2} \sum_{j=1}^N M_j \omega_j^2 R_j^2 + \sum_{j=1}^N g_j R_j$$

$$V_b(\mathbf{R}) = -\varepsilon_0 + \frac{1}{2} \sum_{j=1}^N M_j \omega_j^2 R_j^2 - \sum_{j=1}^N g_j R_j$$

$$V_{ab} = -v_0$$



Full dimensional H-models [$(3N_a-6)D$]

H Model	# states	# parameters	Pros	Cons
SBH	2	$2(3N_a-5)$	Analyt. adiabatic rep.	Low flexibility

SBH: Leggett et al. *Rev Mod Phys* **1987**, 59, 1

Example of dynamics with SBH: Landry; Subotnik. *J Chem Phys* **2011**, 135, 191101

Full dimensional H-models [$(3N_a-6)D$]

H Model	# states	# parameters	Pros	Cons
SBH	2	$2(3N_a-5)$	Analyt. adiabatic rep.	Low flexibility
LVC	N_s	$(3N_a-6)N_s(N_s+1)/2$	Number of states	Parameterization
...				

SBH: Leggett et al. *Rev Mod Phys* **1987**, 59, 1

Example of dynamics with SBH: Landry; Subotnik. *J Chem Phys* **2011**, 135, 191101

LVC: Koppel; Domcke; Cederbaum. *Adv Chem Phys* **1984**, 57, 59

Example of LVC parameterization: Aleotti et al. *J Chem Phys* **2021**, 154, 104106

PES models like Spin-Boson and LVC should be parameterized for each molecule. They are not transferrable.

Their formulation using normal modes makes finding correspondence with real molecules difficult.


Combining these models with hybrid QM/Model approaches is not straightforward.

All these problems are avoided with **Force Field** models.

Force fields

Force-field potential energy surface:

$$E = \sum_{bonds} V_b(b) + \sum_{angles} V_a(\theta) + \sum_{dihedrals} V_d(\phi) + \sum_{i < j} [V_e(R_{ij}) + V_{vdW}(R_{ij})]$$



b - distance between two bonded atoms


θ - angle between three bonded atoms

ϕ - dihedral between four bonded atoms


R_{ij} - distance between two atoms in different molecules or separated by four or more bonds

Bonds and angles are harmonic


$$V_b(b) = K_b(b - b_{eq})^2$$



$$V_a(\theta) = K_\theta(\theta - \theta_{eq})^2$$





Restrict to low temperatures.
No dissociation or bond formation.



Dihedrals follow a periodic potential



$$V_d = \frac{V_n}{2} [1 + \cos(n\phi - \gamma)]$$



Non-bonded interactions have two components

1. Electrostatic (Coulomb)

$$V_e = \frac{q_i q_j}{\epsilon R_{ij}}$$





Non-bonded interactions have two components

$$V_{vdW}(R_{ij})$$


2. Van der Waals
[Lennard-Jones potential]

$$V_{vdW} = \frac{A_{ij}}{R_{ij}^{12}} - \frac{B_{ij}}{R_{ij}^6}$$



Force-field potential energy surface:

$$E = \sum_{bonds} K_b (b - b_{eq})^2 + \sum_{angles} K_\theta (\theta - \theta_{eq})^2 + \sum_{dihedrals} \frac{V_n}{2} [1 + \cos(n\phi - \gamma)] + \sum_{i < j} \left[\frac{A_{ij}}{R_{ij}^{12}} - \frac{B_{ij}}{R_{ij}^6} + \frac{q_i q_j}{\epsilon R_{ij}} \right]$$



$$\sum_{bonds} K_b^{(ca-ca)} (b - b_{eq}^{(ca-ca)})^2 \neq \sum_{bonds} K_b^{(ca-c)} (b - b_{eq}^{(ca-c)})^2$$

AMBER force field: Duan *et al.* *J Comput Chem* **2003**, 24, 1999

```
#####
##          ##
## Atom Type Definitions ##
##          ##
#####
```

atom	1	1	CT	"RCH3 Alkane"	6	12.000	4
atom	2	1	CT	"R2CH2 Alkane"	6	12.000	4
atom	3	1	CT	"R3CH Alkane"	6	12.000	4
atom	4	1	CT	"CH4 Methane"	6	12.000	4
atom	5	1	CT	"R4C Alkane"	6	12.000	4
atom	6	2	HC	"HR Alkane"	1	1.008	1
atom	7	3	CM	"R2C=C Alkene"	6	12.000	3
atom	8	3	CM	"RHC=C Alkene"	6	12.000	3
atom	9	3	CM	"H2C=C Alkene"	6	12.000	3
atom	10	4	HC	"H-C=C Alkene"	1	1.008	1
atom	11	5	CA	"CH Benzene"	6	12.000	3
atom	12	6	HA	"H Benzene"	1	1.008	1
atom	13	5	CA	"C Aromatic Fusion"	6	12.000	3
atom	14	1	CT	"CH3 Toluene"	6	12.000	4
atom	15	1	CT	"CH2 Ethyl Benzene"	6	12.000	4
atom	16	7	OH	"OH Alcohol"	8	15.999	2
atom	17	8	HO	"HO Alcohol"	1	1.008	1
atom	18	2	HC	"CH3 Methanol"	1	1.008	1
atom	19	1	CT	"CH2 & CH3 Alcohol"	6	12.000	4
atom	20	1	CT	"CH i-Pr Alcohol"	6	12.000	4
atom	21	1	CT	"C t-Bu Alcohol"	6	12.000	4
atom	22	1	CT	"CH2 Trifluoroethanol"	6	12.000	4
atom	23	9	CT	"CF3 Trifluoroethanol"	6	12.000	4
atom	24	10	OH	"OH Trifluoroethanol"	8	15.999	2
atom	25	8	HO	"HO Trifluoroethanol"	1	1.008	1

Force-field potential energy surface:

$$E = \sum_{bonds} K_b \left(b - b_{eq} \right)^2 + \sum_{angles} K_\theta \left(\theta - \theta_{eq} \right)^2 + \sum_{dihedrals} \frac{V_n}{2} \left[1 + \cos(\mathbf{n}\phi - \gamma) \right] + \sum_{i < j} \left[\frac{A_{ij}}{R_{ij}^{12}} - \frac{B_{ij}}{R_{ij}^6} + \frac{q_i q_j}{\epsilon R_{ij}} \right]$$

```
#####
##                                     ##
##  Bond Stretching Parameters  ##
##                                     ##
#####
```

bond	1	1	268.0	1.5290
bond	1	2	340.0	1.0900
bond	1	5	317.0	1.5100
bond	1	7	320.0	1.4100
bond	1	9	268.0	1.5290
bond	1	10	320.0	1.4100
bond	1	15	222.0	1.8100
bond	1	17	222.0	1.8100
bond	1	19	382.0	1.4480
bond	1	21	317.0	1.5220
bond	1	23	337.0	1.4490
bond	1	27	317.0	1.5220
bond	1	29	340.0	1.0900
bond	1	30	320.0	1.4100
bond	1	31	375.0	1.4900
bond	1	37	367.0	1.4710
bond	1	39	337.0	1.4630
bond	1	41	317.0	1.4950
bond	1	46	317.0	1.5040

Force fields


- AMBER
- CFF
- CHARMM
- COSMOS-NMR
- CVFF
- ECEPP
- GROMOS
- IFF
- MMFF
- MM2
- OPLS
- QCFF
- UFF
- ...

MD Software

- AMBER
- CHARMM
- COSMOS
- CP2K
- GROMACS
- GROMOS
- LAMMPS
- SPARTAN
- TINKER
- ...

[en.wikipedia.org/wiki/Force_field_\(chemistry\)](https://en.wikipedia.org/wiki/Force_field_(chemistry))

en.wikipedia.org/wiki/Molecular_mechanics




youtu.be/xcMSHy3CqXA

QM/MM PES


Hybrid methods combining QM and MM allows to treat reactive events like bond formation and dissociation.

The system is split into reactive (R) and nonreactive (NR) regions.



Subtractive QM/MM (ONIOM)


$$E_{QM/MM} = V_{MM}(R + NR) + V_{QM}(R) - V_{MM}(R)$$



- ☺ No communication between QM and MM routines
- ☹ Require FF for R region
- ☹ No polarization of R due to NR


Additive QM/MM

$$E_{QM/MM} = V_{QM}(R) + V_{MM}(NR) + V_{QM-MM}(R + NR)$$




$$E_{QM/MM} = V_{QM}(R) + V_{MM}(NR) + V_{QM-MM}(R + NR)$$

$V_{QM}(R)$ gets virtual
link atoms




$$E_{QM/MM} = V_{QM}(R) + \textcolor{red}{V_{MM}}(NR) + V_{QM-MM}(R + NR)$$

MM doesn't need
parameters for R



$$E_{QM/MM} = V_{QM}(R) + V_{MM}(NR) + V_{QM-MM}(R+NR)$$

$V_{QM-MM}(R+NR)$
bonded and
Lennard-Jones
interactions are
handled by FF



$$E_{QM/MM} = V_{QM}(R) + V_{MM}(NR) + V_{QM-MM}(R+NR)$$

Electrostatic interactions in $V_{QM-MM}(R+NR)$ can be FF or QM:

1. Mechanical embedding


$$H_{QM} = H_{QM}(R)$$

$$V_e(R - NR) = \sum_{\substack{i \in R \\ j \in NR}} \frac{q_i q_j}{\epsilon R_{ij}}$$

2. Electrostatic embedding

$$H_{QM} = H_{QM}(R) + \sum_{i \in NR} q_i$$

$$V_e(R - NR) = 0$$



Electrostatic embedding is mandatory for problems involving charge transfer and excited states.

Some advanced aspects in QM/MM

- Certain cases may require polarization of NR too.

This is done with **polarizable force fields**.

Baker. *WIREs: Comp Mol Sci* **2015**, 5, 241

- R and NR regions may change during dynamics.

It is addressed with **adaptive QM/MM**.

Duster *et al.* *WIREs: Comp Mol Sci* **2017**, 7, e1310

- QM/MM input is prone to errors (too many steps).

Input automatization is fundamental.


Pedraza-González *et al.* *J Chem Theory Comput* **2019**, 15, 3134

Software allowing QM/MM

- QM/MM (Truhlar's group)
- ChemShell (Molcas, Turbomole, ...)
- COBRAMM
- CP2K
- Gaussian
- Gamess
- Q-Chem
- NWChem
- ORCA
- MOPAC
- ...

Continuum models


Consider a solvated molecule.



Replace the solvent with a continuum medium.

The molecular charges polarize the medium.


The polarized medium affects the molecular charges.



Continuum (or implicit) solvation models describe the molecular electronic energy changes due to the influence of a polarized continuum medium.


Electrostatic potential V of the polarized medium (V_E) plus the molecule (V_M) is given by the Poisson equation

$$\nabla^2 V = -\frac{4\pi\rho}{\epsilon_E}$$




This problem is equivalent to finding the potential of a fictitious charge distribution σ_E on the surface of the cavity between the molecule and the medium.

$$V_\sigma(\mathbf{r}) = \int_{\Gamma} \frac{\sigma_E(\mathbf{s})}{|\mathbf{r} - \mathbf{s}|} d^2\mathbf{s}$$



Once σ_E is found, the contribution of the solvent to the molecular energy can be computed.

$$G_E = \frac{1}{2} \int_{\Gamma} \sigma_E(s) \left[\int_{vol} \frac{\rho_M(\mathbf{r})}{|\mathbf{r} - s|} d^3 r \right] d^2 s$$




Most implicit models may not describe:


- Charge transfer
- Hydrogen bonds
- Anisotropic and inhomogeneity effects
- Electronic density equilibration after a state change.

They may also have PES discontinuities, which are critical for dynamics.

Charge transfer and discontinuities are well treated with SwiG-PCM.



MS-CASPT2




Li et al. *J Chem Theory Comput* **2015**, 11, 1674


More comparisons: Klamt et al. *J Chem Theory Comput* **2015**, 11, 4220

Photophysics in active environments


(a) Complete QM




(b) Implicit environment



(c) QM/MM



(d) Fragment QM/MM



[Check for updates](#)

Cite this: DOI: 10.1039/d3cp00247k

Surface hopping modeling of charge and energy transfer in active environments

Josene M. Toldo, *^a Mariana T. do Casal, †^a Elizete Ventura, ^b
Silmar A. do Monte *^b and Mario Barbatti *^{ac}

Received 17th January 2023,

Accepted 1st March 2023

DOI: 10.1039/d3cp00247k


rsc.li/pccp

An active environment is any atomic or molecular system changing a chromophore's nonadiabatic dynamics compared to the isolated molecule. The action of the environment on the chromophore occurs by changing the potential energy landscape and triggering new energy and charge flows unavailable in the vacuum. Surface hopping is a mixed quantum-classical approach whose extreme flexibility has made it the primary platform for implementing novel methodologies to investigate the nonadiabatic dynamics of a chromophore in active environments. This Perspective paper surveys the latest developments in the field, focusing on charge and energy transfer processes.


What's an active environment?

An active environment change the photophysics of the chromophore


(a) Isolated chromophore



(b) Passive environment




(c) Active environment



Processes in active environments

- Charge transfer
 - Electron transfer
 - Proton transfer
- Energy transfer
 - Electronic energy
 - Vibrational energy


a) Electron transfer




b) Hydrogen/Proton transfer




c) Electronic energy transfer




d) Internal conversion



e) Intramol. vib. relaxation




f) Vibrational cooling




Charge transfer


Electron and proton transfer processes



H-transfer pathway in adenosine



Electron goes from **GREEN** to **ORANGE**




Be careful with BOMD involving proton transfer.

The large proton delocalization may require a quantum treatment.


Energy transfer

Electronic energy transfer


a) Förster resonance energy transfer (FRET)



b) Dexter energy transfer




c) Coherent transport




Vibrational energy transfer


a) Internal conversion



b) Intramol. vib. redistribution




c) Vibrational cooling




Cytosine in

Argon




25 ps

Benzene



4 ps

Water



1 ps

To know more:

QM/MM

- Groenhof, *In Biomolecular simulations*, **2013**

Active environments

- Toldo et al. *PCCP* **2023**, 25, 8293

Papers available for download at:

amubox.univ-amu.fr/s/xXAiMZrDPb9RMRX (Ask me for the password)